

**DERMATOLOGY +
PLASTIC SURGERY**
For Your Best Self
SALUJA | NOVO | GASGARTH

coolpeel.

ARRIVE 30 MINUTES PRIOR TO APPOINTMENT.

YOU WILL BE AT THE OFFICE FOR APPROXIMATELY 2 HOURS.

PLEASE REPEAT COOLPEEL IF STARTING A SERIES, ANNUALLY FOR MAINTENANCE.

THIS IS ONE OF OUR PATIENTS' FAVORITE CUSTOMIZED SKIN REJUVENATION TREATMENTS!

Please have the following items ready before your treatment day:

- Prescriptions
- Pre and post procedure skin care products

2 Weeks Before Treatment

- Discontinue excessive sun exposure 2 weeks prior to treatment: you should not have sunburn, suntan or spray tan.
- Wear **minimum SPF 30** daily and follow other sun protection measures such as avoiding midday sun and wearing a hat and sunglasses.

1 Week Before Treatment

- If recommended by your practitioner, begin your 4% Hydroquinone cream prescription, Alastin Alluminate, Alastin Skin Regenerating Nectar, and/or Obagi Clear to area twice daily.
- **Discontinue topical retinols, Retin-A/Renova/tretinoin, glycolic or other hydroxy acids.**

1 Day Before Treatment

- If you have a history of cold sores (herpes simplex) Start prescription antiviral medication: **Famvir 500 mg po twice daily x 7 days (Dispense #14).**

Day of Treatment:

- **Please come to the office with a clean face, no makeup, and your hair back away from your face and neck in a band or tie. Wear a loose cut, or buttoned shirt.**
- Remember to finish all of your oral medication as prescribed.
- You will feel warm after treatment. You may take Tylenol 500 mg, two tablets, every 6 hours. Please inform us if there is discomfort not controlled with Tylenol.

What to do: First week

- OK to shower. Wait at least 48 hours to resume mineral makeup.
- Hand-washing is the most important thing you can do to prevent infection.
- Sleep on a clean pillowcase.

6545 N. Wickham Road, Suite F101
Melbourne, FL 32940
321-241-1160
Revised: 1/31/2023

- Apply Nectar to clean skin first (morning and night), followed by a thin layer of Soothe & Protect/ (or Aquaphor) ointment over treated areas during the first 3 to 7 days post procedure
- Do not pick at skin, this is critical to prevent crusting and risk of possible scarring (very rare). Do not rub, scrub or use any exfoliants during this time.

What to do: After First Week and Beyond

- Practice sun avoidance/protection when possible and choose one of the medical grade sunscreens we offer that suits you best - so that you will use it daily. Wearing a hat and sunglasses also helps.
- **Restart your skincare routine.** Do not neglect this! Consider Alastin Alluminate or 4% hydroquinone cream once or twice daily as tolerated if you tend to get brown spots.
Make sure your plan includes:
 - **sunscreen**
 - **topical antioxidant, growth factor, and/or peptide**
 - **brightening serum *if prone to brown spots**
 - **tretinoin cream.**
- It is okay to use makeup.
- If anything is causing increasing redness, discontinue it and inform our office.
- **Aesthetician treatments every 3 months are highly recommended to enhance and maintain your results.**

What you might experience after the procedure:

1. **Heat** may linger for 2-3 hours after the procedure is completed
2. **Redness and swelling** are most common.
3. **Acne or milia formation:** Flare-up of acne or formation of milia possible and may appear 2-4 days post treatment. This is usually self-limiting and resolves within one week. Do not pick at lesions.
4. Risk of **infection** exists despite our best preventative effort. Wash your hands! Keep your environment clean! This is a must.
5. **Itching** usually occurs as a part of the normal wound healing process, but may occur as part of infection, poor wound healing, or contact dermatitis. This usually occurs 3-4 days post procedure and may last up longer. Cold compresses (20 minutes on and off) as needed to alter sensation can help. An oral antihistamine such as Zyrtec or Benadryl can also help. Additional medications should only be used as directed by your practitioner.
6. **Pigmentary changes:** After treatment, rarely, skin may appear darker (hyperpigmentation) or lighter (hypopigmentation) than before.

Call us if you experience any one of the following very rare symptoms or signs:

- Pain that is not relieved by Tylenol. Oozing beyond 48 hours or shallow ulcers. Excessive redness, swelling, bleeding, and/or crust or scab formation. Excessive itching, especially when accompanied by signs of infection: whitish to yellowish film on treated area, with or without foul odor. Fever of 100.4 or above.